

CONVERGENCE OF DISCRETE GREEN FUNCTIONS WITH NEUMANN BOUNDARY CONDITIONS

SHIRSHENDU GANGULY AND YUVAL PERES

ABSTRACT. In this note we prove convergence of Green functions with Neumann boundary conditions for the random walk to their continuous counterparts. Also a few Beurling type hitting estimates are obtained for the random walk on discretizations of smooth domains. These have been used recently in the study of a two dimensional competing aggregation system known as *Competitive Erosion*. Some of the statements appearing in this note are classical for $\mathbb{Z}^2$. However additional arguments are needed for the proofs in the bounded geometry setting.

1. INTRODUCTION

The article considers simple random walk on discretizations of smooth planar simply connected domains. It has two parts. In the first part we show convergence of a class of Green functions with Neumann boundary conditions for the random walk to the corresponding one for Reflected Brownian motion on the same domain. Such convergence results are typically technically challenging and are often extremely useful while understanding scaling limits of statistical physics models. For e.g. results in [10] were used in understanding a modified version of Diffusion Limited Aggregation (DLA). More recently convergence of “discrete analytic functions” to their continuous counterparts were used to show convergence and conformal invariance of well known critical two dimensional statistical physics models. See [15] and the references therein for a detailed account of such results. The results appearing in this article have been used recently in [9] to establish *conformal invariance* of a competing aggregation system known as *Competitive Erosion* on smooth domains. Most of the proofs rely on [5], [7] and [8] where random walk on discretizations of regular domains was shown to converge to Reflected Brownian motion under suitable time change and subsequent local CLT estimates were obtained.

We remark that similar results appear in [10], in the setting of the Dirichlet problem.

In the second part we prove some Beurling type and other hitting estimates for random walk in this setting. Most of the results in this part are classical when the underlying lattice is the whole of $\mathbb{Z}^2$. However for bounded geometry, additional arguments involving heat kernel estimates for the random walk are required.

1.1. Informal set up. Given any simply connected domain $\mathbb{U} \subset \mathbb{C}$ with certain regularity properties, consider its discretization $\mathbb{U}_n := \mathbb{U} \cap \frac{1}{n}\mathbb{Z}^2$. Abusing notation a little we denote by $\mathbb{U}_n$ the graph where the edges are induced by the nearest neighbor edges on $\frac{1}{n}\mathbb{Z}^2$ (formal definition appears later). Fix two points x_1, x_2 on $\mathbb{U}_n$ near the boundary of $\mathbb{U}$ and consider the function on $\mathbb{U}_n$ which is harmonic on all points except x_1 , and x_2 where the discrete laplacian is 1 and -1 respectively. (For the formal definition of laplacian see (3.5)). The goal of this note is to understand convergence of these functions as the mesh size goes to 0. However for technical purposes we study ‘smoothed’ versions of such functions whose laplacian vanishes except on small open sets near the points x_1 and x_2 . More precisely, fix a small $\delta > 0$ and choose the open sets to be balls of radius roughly δ and also at distance δ from x_1 and x_2 , (see Fig 1). We consider the function on $\mathbb{U}_n$ which is harmonic

at every lattice point outside the balls and on the balls the laplacian is roughly the inverse of the the number of lattice points inside the ball. Thus the singularities now are uniformly distributed over open sets in the interior instead of being at points. Formal definition appears in the next section. Such functions were useful in the recent study a competing aggregation system known as *Competitive Erosion*. For more details see [9]. We remark that the proof techniques in this article are general and should work for a much more general class of laplacian conditions.

2. FORMAL DEFINITIONS AND SETUP

$\mathbb{C}$ will denote the complex plane. For any two points $x, y \in \mathbb{C}$, $d(x, y)$ will denote the euclidean distance between them. Also for any set $A \subset \mathbb{C}$ and any $x \in \mathbb{C}$ denote by $d(x, A)$, the distance between the point and the set. $\mathbb{D}$ will be used to denote the unit disc centered at the origin in the complex plane.

Definition 1. For any domain $B \subset \mathbb{C}$ denote by ∂B the boundary of B . Also for any graph $G = (V, E)$ with vertices V and edges E , for any $A \subset V$ let

$$\partial_{out} A := \{y \in A^c : \exists x \in A \text{ such that } x \sim y\}. \quad (2.1)$$

Let a bounded simply connected domain $\mathbb{U} \subset \mathbb{C}$ is “smooth” mean that the boundary of $\mathbb{U}$ is an analytic curve (equivalently the conformal map from $\mathbb{U}$ to $\mathbb{D}$ has a conformal extension across the boundary, see [13, Prop 3.1]). From now on all our domains will be bounded, simply connected and smooth. Hence we will drop the adjectives for brevity.

Setup 1. Given $\mathbb{U}$ we take $\mathbb{U}_n = \mathbb{U} \cap (\frac{1}{n}\mathbb{Z}^2)$, as our vertex set. As the edges of our graph we take the usual nearest-neighbor edges of $\mathbb{U}_n$ though of as a subset of $\frac{1}{n}\mathbb{Z}^2$. However we delete every such edge which intersects $\mathbb{U}^c$. By the smoothness assumption on $\mathbb{U}$, $\mathbb{U}_n$ will be connected for large enough n . See Remark 2.1 below.

Fix $x_1, x_2 \in \partial\mathbb{U}$. For small enough $\delta > 0$ let $y_1, y_2 \in \mathbb{U}$ be such that,

$$\begin{aligned} d(x_i, y_i) &= \delta \\ d(y_i, \partial\mathbb{U}) &> \delta/2. \end{aligned}$$

For $i = 1, 2$, let $\mathbb{U}_i = B(y_i, \frac{\delta}{4})$ (we will call them ‘blobs’). As discrete approximations of $\mathbb{U}_i$ we take

$$\mathbb{U}_{i,n} = B(z_{i,n}, \frac{\delta}{4}) \cap \mathbb{U}_n,$$

where $z_{i,n} \in \frac{1}{n}\mathbb{Z}^2$ is the closest lattice point to y_i .

Note that in the above, y_i ’s were just required to satisfy certain properties and other than that were completely arbitrary. Also we abuse notation a little in the definition of the blobs: n should be thought of as large and hence $\mathbb{U}_n$ (the underlying graph) should not be confused with the blobs $\mathbb{U}_1$ and $\mathbb{U}_2$.

Remark 2.1. The smoothness assumption on $\mathbb{U}$ allows us to choose the y_i ’s. This is formally proved in Corollary 5.1. See Fig. 1. The connectedness of $\mathbb{U}_n$ for large enough n follows due to the locally half plane like behavior, see (5.1).




FIGURE 1. y_i 's are points at distance δ from x_i 's. They are at distance at least $\frac{\delta}{2}$ from $\partial\mathbb{U}$. The blobs are discs of radius $\frac{\delta}{4}$ centered at the y_i 's.

Remark 2.2. For the domain $\mathbb{U}$ and points $x_1, x_2 \in \partial\mathbb{U}$ let

$$\begin{aligned} \phi : \mathbb{D} &\rightarrow \mathbb{U} \\ \psi : \mathbb{U} &\rightarrow \mathbb{D} \end{aligned} \tag{2.2}$$

be conformal maps such that $\phi \circ \psi, \psi \circ \phi$ are the identity maps on the respective domains and $\psi(x_1) = -i, \psi(x_2) = i$. The existence of such maps is guaranteed by the Riemann Mapping Theorem. See for eg: [1, Chapter 6]. In fact there exists a family of such pairs since a conformal map between domains has three degrees of freedom and here we have fixed the value at only two points. However we choose a particular pair (ϕ, ψ) assumed to be fixed throughout the rest of the article. Since $\mathbb{U}$ is smooth, using Schwarz reflection ϕ and hence ψ can be extended conformally across the boundary onto some neighborhoods of $\overline{\mathbb{D}}$ and $\overline{\mathbb{U}}$. In particular this implies that the derivatives $|\phi'|$ and $|\psi'|$ are bounded away from 0 and ∞ on $\overline{\mathbb{D}}$ and $\overline{\mathbb{U}}$ respectively. See [13, Prop 3.1]. This bi-Lipschitz nature of the maps will be used in several distortion estimates throughout the rest of the article. Also note, by construction $|\mathbb{U}_{1,n}| = |\mathbb{U}_{2,n}|$. This will be technically convenient.

2.1. Assumptions, notations and conventions. We summarize some of the notations already used and introduce some new notations and conventions to be used in the sequel.

Through out the article, by random walk on $\mathbb{U}_n$, we will mean the continuous time random walk with $\exp(2n^2)$ waiting times (mean $\frac{1}{2n^2}$) unless specifically mentioned otherwise. This is done to ensure that the random walk density converges to that of Reflected Brownian motion. A fact which would be used heavily.

We will denote the complex plane by $\mathbb{C}$. For any two points $x, y \in \mathbb{C}$, $d(x, y)$ will be used to denote the euclidean distance between them. Also for any set $A \subset \mathbb{C}$ and any $x \in \mathbb{C}$ denote by $d(x, A)$, the distance between the point and the set. $B(x, \epsilon)$ denotes the open euclidean ball of radius ϵ with center x . For any process, and a subset A of the corresponding state space, $\tau(A)$ will denote the hitting time of that set (we drop the dependence on the process in the notation since it will be clear from context). Also $\mathbf{1}(\cdot)$ will be used to denote the indicator function.

To avoid cumbersome notation, we will often use the same letter (generally C, D, c or d) for a constant whose value may change from line to line. $O(\cdot), \Omega(\cdot), \Theta(\cdot)$ are used to denote their usual meaning.

3. PART I: GREEN FUNCTION: DEFINITIONS AND RESULTS

3.1. Discrete Green Function. Recall that we consider continuous time random walk on $\mathbb{U}_n$ with exponential waiting times with mean $\frac{1}{2n^2}$ (Section 2.1). Call it $X(t)$. For $x, y \in \mathbb{U}_n$ let

$$\mathbb{P}_x(X(t) = y) \quad (3.1)$$

denote the chance that the random walk on $\mathbb{U}_n$ starting from x is at y at time t . For notational simplicity we suppress the n dependence in $\mathbb{P}$ since the graph will be clear from context. Similarly for any set $A \subset \mathbb{U}_n$, let $\mathbb{P}_x(X(t) \in A)$ denote the chance that the random walk is in A at time t starting from x .

Definition 2. (Green function) Define the function G_n on $\mathbb{U}_n$: for any $x \in \mathbb{U}_n$,

$$G_n(x) := \frac{2n^2}{|\mathbb{U}_{1,n}|} \int_0^\infty [\mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n})] dt - c. \quad (3.2)$$

where $c = c(\delta)$ is some constant explicitly mentioned in (4.2).

The dependence of G_n on δ (through $\mathbb{U}_{1,n}, \mathbb{U}_{2,n}$) is suppressed in the notation. The centering constant c is not important. The only purpose of the centering is to ensure that for small δ if n is very large then the Green function upto a universal multiplicative constant (domain independent) approaches the function $\log \left| \frac{\psi(x)-i}{\psi(x)+i} \right|$, (where ψ was defined in (2.2)). The convergence results are stated in Section 3.2.

Observe that the Green function upto translation is the difference in the amount of time random walk spends in $\mathbb{U}_{1,n}$ and in $\mathbb{U}_{2,n}$ respectively. It is shown later that the “discrete laplacian” (see (3.5)) of G_n is

$$\frac{1}{|\mathbb{U}_{1,n}|} (\mathbf{1}(\mathbb{U}_{1,n}) - \mathbf{1}(\mathbb{U}_{2,n})).$$

Thus as δ goes to 0, the functions G_n can be thought of as ‘smoothly’ approximating the function on $\mathbb{U}_n$ which is discrete harmonic on $\mathbb{U}_n$ and has laplacian 1 and -1 at x_1 and x_2 respectively. As discussed in the beginning, the purpose of this article is to prove convergence of such ‘smoothed’ Green functions.

The fact that the integral in the expression for G_n is absolutely integrable follows from the following lemma.

Lemma 3.1. *Given $\mathbb{U}$ as in Setup 1 there exists a constant $D = D(\mathbb{U})$ and a time $T = T(\mathbb{U})$ such that, for all large enough n ,*

$$\sup_{x \in \mathbb{U}_n} |\mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n})| \leq 2e^{-Dt}$$

for all $t \geq T$.

Proof. Let us consider the random walk on $\mathbb{U}_n$. The following is a standard consequence of the sub-multiplicative nature of the worst case total variation norm $d_{TV}(\cdot)$:

$$d_{TV}(\ell t_{mix}(1/4)) \leq 2^{-\ell} \quad (3.3)$$

where $t_{mix}(1/4)$ is the $\frac{1}{4}$ total variation mixing time, (see [11, (4.34)]). Let π_{RW} be the stationary measure for the random walk on $\mathbb{U}_n$. It is well known and easy to verify that for any $z \in \mathbb{U}_n$,

$$\pi_{RW}(z) = \frac{d_z}{\sum_{z' \in \mathbb{U}_n} d_{z'}} \quad (3.4)$$

where d_z is the degree of the vertex z in $\mathbb{U}_n$. Recall from Setup 1 that,

$$|\mathbb{U}_{1,n}| = |\mathbb{U}_{2,n}|$$

and also that all points in $\mathbb{U}_{1,n}$ and $\mathbb{U}_{2,n}$ are in the interior of $\mathbb{U}$ and hence have four neighbors for large n . Thus

$$\pi_{RW}(\mathbb{U}_{1,n}) = \pi_{RW}(\mathbb{U}_{2,n}).$$

By definition for any $t > 0$,

$$\sup_{\substack{x \in \mathbb{U}_n \\ i=1,2}} |\mathbb{P}_x(X(t) \in \mathbb{U}_{i,n}) - \pi_{RW}(\mathbb{U}_{i,n})| \leq d_{TV}(t).$$

Thus uniformly over $x \in \mathbb{U}_n$,

$$\begin{aligned} |\mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n})| &\leq |\mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \pi_{RW}(\mathbb{U}_{1,n})| + |\mathbb{P}_x(X(t) \in \mathbb{U}_{2,n}) - \pi_{RW}(\mathbb{U}_{2,n})| \\ &\leq 2d_{TV}(t). \end{aligned}$$

The result now follows from (3.3) and the standard fact that $t_{mix}(1/4) = O(1)$ on $\mathbb{U}_n$, (see Lemma 5.4). $\square$

For any function $f : \mathbb{U}_n \rightarrow \mathbb{R}$ define the laplacian $\Delta f : \mathbb{U}_n \rightarrow \mathbb{R}$, where for any $x \in \mathbb{U}_n$,

$$\Delta f(x) := f(x) - \frac{1}{d_x} \sum_{y \sim x} f(y), \quad (3.5)$$

(d_x is the degree of the vertex x and $y \sim x$ denotes that y is a neighbor of x).

Lemma 3.2. *Consider the function $G_n(\cdot)$ on $\mathbb{U}_n$. Then,*

$$\Delta(G_n) = \frac{1}{|\mathbb{U}_{1,n}|} (\mathbf{1}(\mathbb{U}_{1,n}) - \mathbf{1}(\mathbb{U}_{2,n})), \quad (3.6)$$

where for any subset $A \subset \mathbb{U}_n$, $\mathbf{1}(A)$ denotes the indicator of the set A .

Proof. Proof follows from definition of G_n and looking at the first step of random walk which by definition is of expected duration $\frac{1}{2n^2}$. Thus we have

$$G_n(x) = \frac{1}{|\mathbb{U}_{1,n}|} (\mathbf{1}(\mathbb{U}_{1,n}) - \mathbf{1}(\mathbb{U}_{2,n}))(x) + \frac{1}{d_x} \sum_{y \sim x} G_n(y)$$

and hence the lemma. $\square$

3.2. Main Result. In this section we state the main convergence result of this paper. We start by defining the following function on the domain $\mathbb{U}$, (recall the sets $\mathbb{U}_1, \mathbb{U}_2$ from Setup 1):

$$\tilde{f} := \frac{16}{\text{area}(\mathbb{U}_1)} (\mathbf{1}(\mathbb{U}_2) - \mathbf{1}(\mathbb{U}_1)). \quad (3.7)$$

16 is a constant that falls out of some natural integrals involving the heat kernel of Reflected Brownian motion and is not important. One could normalize things to make the constant 1, however we choose not to do that.

Recall the functions ϕ and ψ from (2.2). Let $\psi(\mathbb{U}_i) =: A_i$. Then,

$$\tilde{f} \circ \phi = \frac{16}{\text{area}(\mathbb{U}_1)} (\mathbf{1}(A_2) - \mathbf{1}(A_1)), \quad (3.8)$$

is a function on $\mathbb{D}$. For any $\mathbb{U}$ as in Setup 1, define the function $G_* : \overline{\mathbb{U}} \rightarrow \mathbb{R}$ such that for all $z \in \overline{\mathbb{U}}$ if $y \in \mathbb{D}$ is such that $\phi(y) = z$, then

$$G_*(z) = \frac{1}{\pi} \int_{|\zeta| < 1} \tilde{f} \circ \phi(\zeta) |\phi'(\zeta)|^2 \log(|\zeta - y|(1 - \bar{\zeta}y)|^2) d\xi d\eta, \quad (3.9)$$

where $\zeta = \xi + i\eta$. Recall that by the smoothness assumption on $\mathbb{U}$ the maps ϕ and ψ have extensions across the boundaries of $\mathbb{D}$ and $\mathbb{U}$ respectively and hence G_* can be defined on $\overline{\mathbb{U}}$. Notice the dependence of G_* on δ through $\tilde{f}$. However for brevity we choose to suppress the dependence on δ (see Setup 1) in the notation.

Even though the function G_n in (3.2) is defined on the graph $\mathbb{U}_n$, to state the next result, we use linear interpolation to think of it as a function on the closure of the whole domain, $\overline{\mathbb{U}}$. We are now ready to state one of the main results of this paper. The interpolation scheme is defined precisely in Subsection 4.3 where the proof of the result appears. Informally it is done in the following way:

- (i). Extend the function G_n from $\mathbb{U}_n$ to $\frac{1}{n}\mathbb{Z}^2$ by fixing it to be 0 outside $\mathbb{U}_n$.
- (ii). Extend the function G_n to all the edges of $\frac{1}{n}\mathbb{Z}^2$ by linearly interpolating the values on the vertices.
- (iii). Extend it to each face of $\frac{1}{n}\mathbb{Z}^2$ so that it is a harmonic function on each face given the value on the edges. Thus the function is extended to the entire complex plane $\mathbb{C}$. By abusing notation a little we still denote the extended function by G_n as well.

Theorem 3.1. *For all small enough δ ,*

$$\lim_{\substack{m \rightarrow \infty \\ n = 2^m}} \sup_{z \in \overline{\mathbb{U}}} |G_n(z) - G_*(z)| = 0.$$

Before proving the above, we remark (see [9, Lemma 5.3]) that as δ goes to 0 the function $G_*(\cdot)$ approaches (up to an explicit multiplicative constant) the function

$$\log \left| \frac{\psi(\cdot) - i}{\psi(\cdot) + i} \right|. \quad (3.10)$$

4. PROOF OF THEOREM 3.1

The proof of Theorem 3.1 involves developing some tools using convergence of random walk on $\mathbb{U}_n$ to Reflected Brownian motion on $\overline{\mathbb{U}}$. For a formal definition of Reflected Brownian motion on $\overline{\mathbb{U}}$ see [7, Definition 2.7]. Also see [2, 6, 5]. Throughout the rest of the article we will denote it by B_t . Consider the function, for $z \in \overline{\mathbb{U}}$,

$$G(z) := \frac{2}{\text{area}(\mathbb{U}_1)} \int_0^\infty [P_z(B_t \in \mathbb{U}_1) - P_z(B_t \in \mathbb{U}_2)] dt - c. \quad (4.1)$$

where for $i = 1, 2$, $P_z(B_t \in \mathbb{U}_i)$ denotes the probability that started from z , B_t is in $\mathbb{U}_i$. The constant c is chosen such that the integral of G along $\partial\mathbb{U}$ is 0 where we parametrize the boundary $\partial\mathbb{U}$ by $\theta \in [0, 2\pi)$ via the conformal map ϕ (2.2). Formally we fix c such that

$$\int_{|\zeta|=1} G \circ \phi(\zeta) \frac{d\zeta}{\zeta} = 0. \quad (4.2)$$

Compare the expression of $G(\cdot)$ with $G_n(\cdot)$ defined in (3.2) (note that the constant c is the same in both expressions).

Before providing formal arguments we sketch the general outline of the proof first. The proof of Theorem 3.1 has two parts: in the first part we show that G_n converges to G . This will follow by

convergence of the random walk measure on $\mathbb{U}_n$ to Reflected Brownian motion B_t . For more on this see [5], [7] and the references therein. Thus the only remaining step then is to show that indeed

$$G_* = G.$$

This will be proved using the fact that the density for Reflected Brownian motion is a fundamental solution to the Neumann problem and hence the function G roughly satisfies,

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) G = \frac{4}{\text{area}(\mathbb{U}_1)} (\mathbf{1}(\mathbb{U}_2) - \mathbf{1}(\mathbb{U}_1)) \quad (4.3)$$

with Neumann boundary conditions. One then checks that G_* is a solution to the above Neumann problem as well. The proof is then complete by uniqueness of the solution of such a problem which allows us to conclude that $G_* = G$. We adopt the following standard notation:

$$\Delta := \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}. \quad (4.4)$$

Recall (3.5). Thus we use Δ to denote the laplacian in both the continuous and discrete setting since there will be no scope of confusion.

4.1. Continuum version of G_n . We begin by studying the function $G(z)$ defined in (4.1). Let B_t be Reflected Brownian motion (RBM) on $\overline{\mathbb{U}}$ and $p(t, x, y)$ be the heat kernel of B_t defined on

$$\mathbb{R}_+ \times \overline{\mathbb{U}} \times \overline{\mathbb{U}},$$

i.e. $p(t, x, y)$ is the density of RBM started from x at time t at point y . Before proceeding we state some classical results about regularity properties of $p(t, x, y)$.

Theorem 4.1.

- a. [14, Lemma 2.1] $p(t, x, y)$ is continuous on $(0, \infty) \times \overline{\mathbb{U}} \times \overline{\mathbb{U}}$.
- b. [14, Theorem 2.1] Let f be a compactly supported C^∞ function defined on $\mathbb{U}$. Then

$$\int_{\overline{\mathbb{U}}} p(t, x, y) f(y) dy$$

is continuous on $(0, \infty) \times \overline{\mathbb{U}}$.

- c. [14, Theorem 2.2] Let f be a compactly supported C^∞ function defined on $\mathbb{U}$. Then

$$u(t, x) = \int_0^t ds \int_{\overline{\mathbb{U}}} p(s, x, y) f(y) dy$$

has the following properties :

- i. is continuous on $(0, \infty) \times \overline{\mathbb{U}}$, continuously differentiable in t in $(0, \infty)$ and of class $C^2(\mathbb{U})$ and $C^1(\overline{\mathbb{U}})$ as a function of x ,
- ii. $(\frac{\partial}{\partial t} - \frac{1}{2}\Delta) u(t, x) = f(x)$,
- iii. $\frac{\partial}{\partial \nu} u(t, x) = 0$, where $\frac{\partial}{\partial \nu}$ denotes the normal derivative and $x \in \partial\mathbb{U}$,
- iv. $\lim_{t \rightarrow 0} u(t, x) = 0$ uniformly on $\overline{\mathbb{U}}$.

The results quoted from [14] are actually proved in much more generality. However for our purposes the above versions would suffice. For $i = 1, 2$, recall $\mathbb{U}_i$ from Setup 1 and let,

$$p(t, x, \mathbb{U}_i) := \int_{\mathbb{U}_i} p(t, x, y) dy. \quad (4.5)$$

Thus the expression in (4.1) is the same as,

$$G(z) = \frac{2}{\text{area}(\mathbb{U}_1)} \int_0^\infty [p(t, z, \mathbb{U}_1) - p(t, z, \mathbb{U}_2)] dt - c. \quad (4.6)$$

Recall that the above expression depends on δ (Setup 1) which determines the sets $\mathbb{U}_1$ and $\mathbb{U}_2$. However we will suppress the dependence on δ for notational brevity since there is little chance for confusion.

Remark 4.1. The fact that the above integral is absolutely convergent directly follows from the following mixing lemma and the fact that by choice $\text{area}(\mathbb{U}_1) = \text{area}(\mathbb{U}_2)$.

Lemma 4.1. [3, Theorem 2.4] *There exists constants $c = c(\mathbb{U}), T = T(\mathbb{U}) > 0$ such that for all $x, y \in \overline{\mathbb{U}}$ and $t > T$,*

$$|p(t, x, y) - \frac{1}{\text{area}(\mathbb{U})}| \leq e^{-ct}.$$

Note that to be able to use Theorem 4.1 to prove (4.3) one has to approximate the indicator functions on $\mathbb{U}_1, \mathbb{U}_2$ by C^∞ functions. Let $g_1, g_2 \dots$ be C^∞ functions taking values in the interval $[0, 1]$ such that for any $j \in \mathbb{N}$ and $z \in \mathbb{C}$

$$g_j(z) = \begin{cases} 1 & |z| \leq 1 - \frac{1}{j} \\ 0 & |z| \geq 1 - \frac{1}{2j}. \end{cases}$$

That is g_j 's form a sequence of smooth functions approximating from below the indicator function on the unit ball. For any $a \in \mathbb{C}$, $b > 0$ and $j \in \mathbb{N}$ we denote by $g_j(a, b, \cdot)$, the function such that for any $z \in \mathbb{C}$

$$g_j(a, b, z) = g_j\left(\frac{z - a}{b}\right),$$

i.e. $g_j(a, b, \cdot)$ approximates the indicator function on the ball $B(a, b)$. Recall from Setup 1 that $\mathbb{U}_1, \mathbb{U}_2$ have centers y_1, y_2 and radius $\frac{\delta}{4}$. For brevity let $\tilde{\delta} := \frac{\delta}{4}$. The next easy lemma uses the g_j 's to approximate the integrals appearing in (4.6).

Lemma 4.2. *Given any T for every $\epsilon > 0$ there exists J such that for all $j > J$ and $i = 1, 2$*

$$\sup_{z \in \overline{\mathbb{U}}} \left| \int_0^T \left[\int_{\overline{\mathbb{U}}} p(t, z, \zeta) g_j(y_i, \tilde{\delta}, \zeta) d\xi d\eta - p(t, z, \mathbb{U}_i) \right] dt \right| \leq \epsilon$$

where $\zeta = \xi + i\eta$.

Proof. We prove it only for the case $i = 1$. Using the trivial observation that for all $t > 0$, both

$$p(t, z, \mathbb{U}_1) \text{ and } \int_{\overline{\mathbb{U}}} p(t, z, \zeta) g_j(y_1, \tilde{\delta}, \zeta) d\xi d\eta \leq 1,$$

and $p(t, z, \mathbb{U}_1) = \int_{\mathbb{U}} p(t, z, \zeta) \mathbf{1}(\zeta \in \mathbb{U}_1) d\xi d\eta$ it suffices to show ,

$$\sup_{z \in \overline{\mathbb{U}}} \left| \int_{\epsilon/2}^T \left[\int_{\overline{\mathbb{U}}} p(t, z, \zeta) \left(g_j(y_1, \tilde{\delta}, \zeta) - \mathbf{1}(\zeta \in \mathbb{U}_1) \right) d\xi d\eta \right] dt \right| \leq \frac{\epsilon}{2}.$$

Now by Lemma 4.1, $p(t, z, \zeta)$ is bounded on the cylinder $[\epsilon/2, T] \times \overline{\mathbb{U}} \times \overline{\mathbb{U}}$. Also clearly,

$$\lim_{j \rightarrow \infty} \int_{\overline{\mathbb{U}}} |g_j(y_1, \tilde{\delta}, \zeta) - \mathbf{1}(\zeta \in \mathbb{U}_1)| d\xi d\eta = 0. \quad (4.7)$$

Thus we are done. $\square$

In the next couple of lemmas we approximate the Green function (4.6) in terms of the functions $g_j(\cdot, \cdot, \cdot)$.

Lemma 4.3. *For $i = 1, 2$, given $\epsilon > 0$, there exists $T = T(\epsilon)$ such that for all $t \geq T$ and all large enough j ,*

$$\sup_{z \in \bar{\mathbb{U}}} \left| \int_{\bar{\mathbb{U}}} p(t, z, \zeta) g_j(y_i, \tilde{\delta}, \zeta) d\xi d\eta - \frac{\text{area}(\mathbb{U}_1)}{\text{area}(\mathbb{U})} \right| \leq \epsilon,$$

where $\zeta = \xi + i\eta$.

Proof. By Lemma 4.1 for large t and all j ,

$$\sup_{z \in \bar{\mathbb{U}}} \int_{\bar{\mathbb{U}}} \left| p(t, z, \zeta) - \frac{1}{\text{area}(\mathbb{U})} \right| g_j(y_i, \tilde{\delta}, \zeta) d\xi d\eta = O(e^{-ct}). \quad (4.8)$$

Now by (4.7) for large j ,

$$\frac{1}{\text{area}(\mathbb{U})} \left| \int_{\bar{\mathbb{U}}} g_j(y_i, \tilde{\delta}, \zeta) d\xi d\eta - \text{area}(\mathbb{U}_1) \right| \leq \epsilon.$$

Thus the above two statements along with triangle inequality complete the proof. $\square$

Given $j \in \mathbb{N}$ and $T > 0$ define,

$$G_{j,T}(z) = \frac{2}{\text{area}(\mathbb{U}_1)} \int_0^T dt \left[\int_{\bar{\mathbb{U}}} p(t, z, \zeta) [g_j(y_1, \tilde{\delta}, \zeta) - g_j(y_2, \tilde{\delta}, \zeta)] d\xi d\eta \right] - c, \quad (4.9)$$

where c is the same as in (4.1) and $\zeta = \xi + i\eta$. We then have the following lemma showing that $G_{j,T}$ approximate G as j, T go to infinity.

Lemma 4.4.

$$\lim_{j, T \rightarrow \infty} \sup_{z \in \bar{\mathbb{U}}} |G_{j,T}(z) - G(z)| = 0.$$

Proof. Clearly by definition

$$\int_{\mathbb{U}} [g_j(y_1, \tilde{\delta}, \zeta) - g_j(y_2, \tilde{\delta}, \zeta)] d\xi d\eta = 0.$$

Hence for every j and all intervals $[T, T']$ where T is large enough and $T < T'$ by (4.8)

$$\int_T^{T'} dt \left| \int_{\bar{\mathbb{U}}} p(t, z, x) [g_j(y_1, \tilde{\delta}, x) - g_j(y_2, \tilde{\delta}, x)] dx \right| = O(e^{-cT}). \quad (4.10)$$

Also note that by Lemma 4.1 for large enough $T < T'$,

$$\int_T^{T'} |p(t, z, \mathbb{U}_1) - p(t, z, \mathbb{U}_2)| dt = O(e^{-cT}). \quad (4.11)$$

Hence given $\epsilon > 0$, choose T such that both the above integrals in (4.10) and (4.11) are less than ϵ for all $T' > T$ and $z \in \bar{\mathbb{U}}$. The lemma now follows by Lemma 4.2. $\square$

Since g_j 's are compactly supported C^∞ functions, for any j and z using Theorem 4.1 b., c., and fundamental theorem of calculus we have,

$$\frac{\partial}{\partial t} G_{j,t}(z) = \frac{2}{\text{area}(\mathbb{U}_1)} \int_{\bar{\mathbb{U}}} p(t, z, x) [g_j(y_1, \tilde{\delta}, x) - g_j(y_2, \tilde{\delta}, x)] dx. \quad (4.12)$$

Also by Theorem 4.1 c.ii, iii. for any $t > 0$

$$\begin{aligned}\Delta G_{j,t}(z) &= \frac{4}{\text{area}(\mathbb{U}_1)}[g_j(y_2, \tilde{\delta}, z) - g_j(y_1, \tilde{\delta}, z)] + \frac{4}{\text{area}(\mathbb{U}_1)} \int_{\overline{\mathbb{U}}} p(t, z, x)[g_j(y_1, \tilde{\delta}, x) - g_j(y_2, \tilde{\delta}, x)] dx \\ \frac{\partial}{\partial \nu} G_{j,t}(z) &= 0 \text{ for all } z \in \partial \mathbb{U}.\end{aligned}\tag{4.13}$$

4.2. Closed form of the limit. In this subsection we show that the function G is same as the function G_* (3.9).

Theorem 4.2. *Let $\mathbb{U}$ be as in Setup 1. For all $z \in \overline{\mathbb{U}}$,*

$$G(z) = G_*(z).\tag{4.14}$$

To show this, we identify G as a solution to a second order differential equation also satisfied by G_* . The result then follows by uniqueness of such a solution. We first quote a result in the theory of boundary value problems with Neumann boundary condition.

Lemma 4.5. [4, Theorem 8] *A function $w \in C^2(\mathbb{D}) \cap C^1(\overline{\mathbb{D}})$ on the disc satisfying the following properties*

$$\begin{aligned}\Delta(w) &= \frac{f}{4} \\ \frac{\partial w}{\partial \nu} &= \gamma \text{ on } \partial \mathbb{D} \\ \frac{1}{2\pi i} \int_{\partial \mathbb{D}} \frac{w(\zeta)}{\zeta} d\zeta &= d\end{aligned}$$

where $f \in L^1(\mathbb{R}, \mathbb{D})$, $\gamma \in C(\mathbb{R}, \partial \mathbb{D})$ and $d \in \mathbb{R}$ exists iff

$$\frac{1}{2\pi i} \int_{|\zeta|=1} \gamma(\zeta) \frac{d\zeta}{\zeta} = \frac{2}{\pi} \int_{|\zeta|<1} f(\zeta) d\xi d\eta.$$

The unique solution in that case is given by the following

$$w(z) = d - \frac{1}{2\pi i} \int_{|\zeta|=1} \gamma(\zeta) \log(|\zeta - z|^2) \frac{d\zeta}{\zeta} + \frac{1}{\pi} \int_{|\zeta|<1} f(\zeta) \log(|(\zeta - z)(1 - \bar{\zeta}z)|^2) d\xi d\eta,$$

where $\zeta = \xi + i\eta$.

Recall the maps ϕ and ψ from (2.2). They will be used throughout the rest of the subsection. The next simple lemma shows how the laplacian changes under a change of variable.

Lemma 4.6. *For a function $u \in C^2(\mathbb{U}) \cap C^1(\overline{\mathbb{U}})$ with the following properties*

$$\begin{aligned}\Delta(u) &= f \text{ on } \mathbb{U} \\ \frac{\partial u}{\partial \nu} &= 0 \text{ on } \partial \mathbb{U}\end{aligned}$$

the function $v = u \circ \phi$ on $\overline{\mathbb{D}}$ satisfies the following properties,

i. $v \in C^2(\mathbb{D}) \cap C^1(\overline{\mathbb{D}})$.

ii. $\Delta(v) = (f \circ \phi)|\phi'|^2$.

iii. $\frac{\partial v}{\partial \nu} = 0$ on $\partial \mathbb{D}$.

Proof. *i.* is obvious since it is the composition with a conformal map which is analytic across the boundary by hypothesis.

ii. follows from chain rule for differentiation and using the Cauchy-Riemann equations.

iii. Since conformal maps preserve angles it follows that

$$\frac{\partial v}{\partial \nu} = \frac{\partial u}{\partial \nu} |\phi'(z)|.$$

Now by hypothesis

$$\frac{\partial u}{\partial \nu} \Big|_{\partial \mathbb{U}} = 0$$

and hence we are done. $\square$

Using the above observation we state a lemma analogous to Lemma 4.5 for $\mathbb{U}$.

Lemma 4.7. *Let function $w \in C^2(\mathbb{U}) \cap C^1(\bar{\mathbb{U}})$ be a function on $\mathbb{U}$ satisfying the following properties*

$$\begin{aligned} \Delta(w) &= \frac{f}{4} \\ \frac{\partial w}{\partial \nu} &= 0 \text{ on } \partial \mathbb{U} \end{aligned}$$

where $f \in L^1(\mathbb{R}, \mathbb{U})$ then

$$w \circ \phi(z) = d + \frac{1}{\pi} \int_{|\zeta| < 1} f \circ \phi(\zeta) |\phi'(\zeta)|^2 \log(|(\zeta - z)(1 - \bar{\zeta}z)|^2) d\xi d\eta, \quad (4.15)$$

where

$$\frac{1}{2\pi i} \int_{\partial \mathbb{D}} \frac{w \circ \phi(\zeta)}{\zeta} d\zeta = d.$$

Proof. The proof follows from Lemmas 4.5 and 4.6. $\square$

We are now ready to prove Theorem 4.2.

Proof of Theorem 4.2. For any j and T recall $G_{j,T}$ from (4.9). By (4.13), $G_{j,T}$ satisfies the hypotheses of Lemma 4.7 with $\Delta G_{j,T} = \frac{f_{j,T}}{4}$, where

$$f_{j,T}(\cdot) = \frac{16}{\text{area}(\mathbb{U}_1)} \left[g_j(y_2, \tilde{\delta}, \cdot) - g_j(y_1, \tilde{\delta}, \cdot) + \int_{\mathbb{U}} p(T, \cdot, \zeta) [g_j(y_1, \tilde{\delta}, \zeta) - g_j(y_2, \tilde{\delta}, \zeta)] d\xi d\eta \right], \quad (4.16)$$

where $\zeta = \xi + i\eta$. Thus by Lemma 4.7 for all $z \in \bar{\mathbb{D}}$,

$$G_{j,T} \circ \phi(z) = c_{j,T} - c + \frac{1}{\pi} \int_{|\zeta| < 1} |\phi'(\zeta)|^2 (f_{j,T} \circ \phi)(\zeta) \log(|(\zeta - z)(1 - \bar{\zeta}z)|^2) d\xi d\eta, \quad (4.17)$$

where

$$c_{j,T} = \frac{1}{2\pi i} \int_{|\zeta|=1} (G_{j,T} \circ \phi + c) \frac{d\zeta}{\zeta}. \quad (4.18)$$

Recall from (3.7),

$$\tilde{f} = \frac{16}{\text{area}(\mathbb{U}_1)} (\mathbf{1}(\mathbb{U}_2) - \mathbf{1}(\mathbb{U}_1)).$$

and that $\psi(\mathbb{U}_i) = A_i$. Then

$$\tilde{f} \circ \phi = \frac{16}{\text{area}(\mathbb{U}_1)} (\mathbf{1}(A_2) - \mathbf{1}(A_1)).$$

We first prove that

$$\lim_{j,T \rightarrow \infty} \sup_{z \in \mathbb{D}} |(G_{j,T} \circ \phi(z) - c_{j,T} + c) - G_* \circ \phi(z)| = 0. \quad (4.19)$$

Using the expression in (4.17) and (3.9) we get that given $\epsilon > 0$, for large enough j, T , for all $z \in \overline{\mathbb{D}}$

$$\begin{aligned} |G_{j,T} \circ \phi(z) - c_{j,T} + c - G_* \circ \phi(z)| &\leq \frac{32}{\text{area}(\mathbb{U}_1)} \left[\int_{\zeta \in A_1 \setminus \psi(B(y_1, \tilde{\delta}(1-\frac{1}{j})))} |\phi'|^2 |\log(|\zeta - z||1 - \bar{\zeta}z|)| d\xi d\eta \right. \\ &\quad + \int_{\zeta \in A_2 \setminus \psi(B(y_2, \tilde{\delta}(1-\frac{1}{j})))} |\phi'|^2 |\log(|\zeta - z||1 - \bar{\zeta}z|)| d\xi d\eta \\ &\quad \left. + \int_{\zeta \in \mathbb{D}} \epsilon |\phi'|^2 |\log(|\zeta - z||1 - \bar{\zeta}z|)| d\xi d\eta \right]. \end{aligned}$$

The terms on the RHS corresponds to difference between the functions $f_{j,T} \circ \phi$ and $\tilde{f} \circ \phi$. The first term corresponds to difference between $g_j(y_1, \tilde{\delta}, \phi(\cdot))$ and $1(A_1)$. Similarly the second term corresponds to $g_j(y_2, \tilde{\delta}, \phi(\cdot))$ and $1(A_2)$. The last term corresponds to $\int_{\mathbb{U}} p(T, \cdot, \zeta) [g_j(y_1, \tilde{\delta}, \zeta) - g_j(y_2, \tilde{\delta}, \zeta)] d\xi d\eta$. By Lemma 4.3 for any ϵ , for large enough T and j ,

$$\sup_{z \in \overline{\mathbb{U}}} \left| \int_{\mathbb{U}} p(T, z, \zeta) [g_j(y_1, \tilde{\delta}, x) - g_j(y_2, \tilde{\delta}, \zeta)] d\xi d\eta \right| \leq \epsilon.$$

Now in all the above terms we can ignore $|\phi'|^2$ since it is bounded by Remark 2.2. As $\log(|\zeta - z||1 - \bar{\zeta}z|)$ is a uniformly locally integrable function in z we see that by making ϵ small and j large we can make the above quantity arbitrarily small uniformly over $z \in \overline{\mathbb{D}}$. Thus

$$\lim_{j,T \rightarrow \infty} \left[\sup_{z \in \overline{\mathbb{D}}} |G_{j,T}(z) - c_{j,T} + c - G_*(z)| \right] = 0.$$

Now notice by Lemma 4.4

$$\lim_{j,T \rightarrow \infty} \frac{1}{2\pi i} \int_{|\zeta|=1} (G_{j,T} \circ \phi) \frac{d\zeta}{\zeta} = \frac{1}{2\pi i} \int_{|\zeta|=1} (G \circ \phi) \frac{d\zeta}{\zeta}.$$

By the choice of c in (4.2)

$$\frac{1}{2\pi i} \int_{|\zeta|=1} (G \circ \phi) \frac{d\zeta}{\zeta} = 0.$$

Hence by (4.18)

$$\lim_{j,T \rightarrow \infty} c_{j,T} = c.$$

This along with (4.19) completes the proof. □

Remark 4.2. Recall the constant c in (4.1) such that.

$$\int_{|\zeta|=1} G \circ \phi(\zeta) \frac{d\zeta}{\zeta} = 0.$$

Consider the special case of the unit disc with point $x_1 = -i, x_2 = i$ and $y_1 = -(1-\delta)i, y_2 = (1-\delta)i$. Now owing to the invariance of Reflected Brownian motion on $\mathbb{D}$ under the transformation $\zeta \rightarrow -\zeta$, one sees that the integral in (4.1)

$$\frac{2}{\text{area}(\mathbb{D}_1)} \int_0^\infty [P_z(B_t \in \mathbb{D}_1) - P_z(B_t \in \mathbb{D}_2)] dt$$

is an odd function, (since by the symmetric choice of y_1, y_2 , $\mathbb{D}_1 = -\mathbb{D}_2$). Thus c is 0 and

$$G(z) = \frac{2}{\text{area}(\mathbb{D}_1)} \int_0^\infty [P_z(B_t \in \mathbb{D}_1) - P_z(B_t \in \mathbb{D}_2)] dt.$$

4.3. Convergence. In this section we prove the last technical piece needed to complete the proof of Theorem 3.1. Namely we show that the function G_n converges to G . This along with Theorem 4.2 then would clearly complete the proof of Theorem 3.1.

As mentioned in the statement of Theorem 3.1 the function G_n is interpolated from the graph $\mathbb{U}_n$ to $\overline{\mathbb{U}}$. We define precisely the interpolation method which is sketched right before the statement of the theorem. We follow the scheme mentioned in [7, Pf of Theorem 2.12] (also appears in [8, Pf of Theorem 2.2.8]).

For $x, y \in \mathbb{U}_n$, define,

$$p_n(t, x, y) = \frac{\mathbb{P}_x(X(t) = y)}{m_n(y)} \quad (4.20)$$

where $\mathbb{P}_x(X(t) = y)$ is the probability that the continuous time random walk $X(t)$ started from x is at y at time t (see (3.1)). $m_n(y) = \frac{d_y}{4n^2}$ is the symmetrizing measure of the random walk (d_y is the degree of the vertex y).

For all pairs $x, y \in \frac{1}{n}\mathbb{Z}^2$, such that at least one of them is not in $\mathbb{U}_n$ define for all t , $p_n(t, x, y) = 0$. Now having defined $p_n(t, x, y)$ for all $t \in \mathbb{R}_+$, $x, y \in \frac{1}{n}\mathbb{Z}^2$ we extend it to $\mathbb{R}_+ \times \overline{\mathbb{U}} \times \overline{\mathbb{U}}$. To this end we interpolate $p_n(t, x, y)$ by a sequence of harmonic extensions along simplices,

- i.* First harmonically extend it along the edges of $\frac{1}{n}\mathbb{Z}^2$ using the value on the vertices.
 - ii.* Then harmonically extend it to the squares (faces of $\frac{1}{n}\mathbb{Z}^2$) using the value on the edges.
- Thus we have extended $p_n(t, x, y)$ to $\mathbb{R}_+ \times \mathbb{C} \times \mathbb{C}$, and hence in particular to $\mathbb{R}_+ \times \overline{\mathbb{U}} \times \overline{\mathbb{U}}$.

Note that by (4.20) for $x \in \mathbb{U}_n$, the expression of $G_n(x)$ in (3.2) is the same as

$$\frac{2n^2}{|\mathbb{U}_{1,n}|} \int_0^\infty \left[\sum_{y \in \mathbb{U}_{1,n}} \frac{p_n(t, x, y)}{n^2} - \sum_{y \in \mathbb{U}_{2,n}} \frac{p_n(t, x, y)}{n^2} \right] dt - c. \quad (4.21)$$

$p_n(t, x, y)$ is divided by n^2 in the above sums since all $y \in \mathbb{U}_{1,n} \cup \mathbb{U}_{2,n}$ have degree 4.

Now by the above extension $p_n(t, x, y)$ is extended to all $x \in \overline{\mathbb{U}}, y \in \mathbb{U}_{1,n} \cup \mathbb{U}_{2,n}$. The above expression thus allows us to define the extended function $G_n(x)$ on $\overline{\mathbb{U}}$. It is easy to see that since the integral in (4.21) is finite for all $x \in \mathbb{U}_n$, the interpolation does not cause additional issues.

For notational unification we formally denote the two sums inside the integral in (4.21) as $\mathbb{P}_x(X(t) \in \mathbb{U}_{i,n})$ for $i = 1, 2$ respectively as in (3.2) even though now the function lives on $\overline{\mathbb{U}}$ and hence x might not be in $\mathbb{U}_n$.

We now state the following lemma which along with Theorem 4.2 proves Theorem 3.1.

Lemma 4.8.

$$\lim_{\substack{m \rightarrow \infty \\ n=2^m}} \sup_{x \in \overline{\mathbb{U}}} |G_n(x) - G(x)| = 0$$

where G is defined in (4.1).

Before proving the lemma we state a local CLT result for random walk approximation of Reflected Brownian motion.

Theorem 4.3. [7, Theorem 2.12] *For all positive numbers $a < b$,*

$$\lim_{\substack{m \rightarrow \infty \\ n=2^m}} \sup_{a \leq t \leq b} \sup_{x, y \in \bar{\mathbb{U}}} |p_n(t, x, y) - p(t, x, y)| = 0$$

where $p(t, x, y)$ is the heat kernel of Reflected Brownian motion as defined in Subsection 4.1.

Also for any set $A \subset \bar{\mathbb{U}}$ and $x \in \bar{\mathbb{U}}$ let

$$p(t, x, A) := \int_A p(t, x, y) dy.$$

Proof of Lemma 4.8. Fix a small positive number a and a large number T . We split G_n in the following way

$$\begin{aligned} G_n(x) &= \frac{2n^2}{|\mathbb{U}_{1,n}|} \int_0^a \mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n}) dt \\ &+ \frac{2n^2}{|\mathbb{U}_{1,n}|} \int_a^T \mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n}) dt \\ &+ \frac{2n^2}{|\mathbb{U}_{1,n}|} \int_T^\infty \mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n}) dt \end{aligned}$$

and similarly,

$$\begin{aligned} G(x) &= \frac{2}{\text{area}(\mathbb{U}_1)} \int_0^a p(t, x, \mathbb{U}_1) - p(t, x, \mathbb{U}_2) dt \\ &+ \frac{2}{\text{area}(\mathbb{U}_1)} \int_a^T p(t, x, \mathbb{U}_1) - p(t, x, \mathbb{U}_2) dt \\ &+ \frac{2}{\text{area}(\mathbb{U}_1)} \int_T^\infty p(t, x, \mathbb{U}_1) - p(t, x, \mathbb{U}_2) dt. \end{aligned}$$

To prove the lemma we will show that the three terms in the RHS for $G_n(x)$ are close to the corresponding terms for $G(x)$. From Lemma 4.1 it follows that uniformly over $x \in \bar{\mathbb{U}}$ and $t \geq T$

$$|p(t, x, \mathbb{U}_1) - \frac{\text{area}(\mathbb{U}_1)}{\text{area}(\mathbb{U})}| \leq e^{-ct}$$

and similarly for $\mathbb{U}_2$. Now since by hypothesis $\text{area}(\mathbb{U}_1) = \text{area}(\mathbb{U}_2)$,

$$\sup_{x \in \bar{\mathbb{U}}} \int_T^\infty |p(t, x, \mathbb{U}_1) - p(t, x, \mathbb{U}_2)| dt \leq 2 \int_T^\infty e^{-ct} dt.$$

Also by Lemma 3.1 and our interpolation scheme

$$\sup_{x \in \bar{\mathbb{U}}} \int_T^\infty |\mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n})| dt \leq 2 \int_T^\infty e^{-Ct} dt$$

for T large enough. Thus in both the expressions the third term can be made arbitrarily small by choosing T large enough. Now the integral in the first term is at most a and hence can be made small by choosing a small enough. We now show that for any fixed a and T the middle term goes to zero as n goes to infinity. First notice that since $|\mathbb{U}_{1,n}| = n^2 \text{area}(\mathbb{U}_1) + O(n)$ it suffices to just show,

$$\sup_{x \in \bar{\mathbb{U}}} \left| \int_a^T [\mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n})] dt - \int_a^T [p(t, x, \mathbb{U}_1) - p(t, x, \mathbb{U}_2)] dt \right|$$

goes to 0 as $n = 2^m \rightarrow \infty$. Using Theorem 4.3 we choose n large enough such that

$$\sup_{[a,T]} \sup_{x,y \in \bar{\mathbb{U}}} |p_n(t, x, y) - p(t, x, y)| \leq \epsilon$$

for some small number ϵ . Now the continuity of the heat kernel $p(t, x, y)$ allows us to use Riemann sums to approximate

$$\int_a^T [p(t, x, \mathbb{U}_1) - p(t, x, \mathbb{U}_2)] dt.$$

That is

$$\int_a^T (p(t, x, \mathbb{U}_1) - p(t, x, \mathbb{U}_2)) dt = \lim_{n \rightarrow \infty} \int_a^T \left[\sum_{y \in \mathbb{U}_{1,n}} \frac{p(t, x, y)}{n^2} - \sum_{y \in \mathbb{U}_{2,n}} \frac{p(t, x, y)}{n^2} \right] dt. \quad (4.22)$$

More over the above convergence is uniform in x on $\bar{\mathbb{U}}$. This is because $p(t, x, y)$ is continuous on the compact set $[a, T] \times \bar{\mathbb{U}} \times \bar{\mathbb{U}}$. Now using Theorem 4.3 we get that for any fixed a and T

$$\lim_{\substack{n=2^m \\ m \rightarrow \infty}} \sup_{x \in \bar{\mathbb{U}}} \left| \left(\int_a^T \left[\sum_{y \in \mathbb{U}_{1,n}} \frac{p_n(t, x, y)}{n^2} - \sum_{y \in \mathbb{U}_{2,n}} \frac{p_n(t, x, y)}{n^2} \right] dt \right) - \left(\int_a^T \left[\sum_{y \in \mathbb{U}_{1,n}} \frac{p(t, x, y)}{n^2} - \sum_{y \in \mathbb{U}_{2,n}} \frac{p(t, x, y)}{n^2} \right] dt \right) \right| = 0.$$

Also by definition for all $x \in \bar{\mathbb{U}}$,

$$\int_a^T \left[\sum_{y \in \mathbb{U}_{1,n}} \frac{p_n(t, x, y)}{n^2} - \sum_{y \in \mathbb{U}_{2,n}} \frac{p_n(t, x, y)}{n^2} \right] dt = \int_a^T [\mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n})] dt.$$

Thus by (4.22)

$$\lim_{\substack{n=2^m \\ m \rightarrow \infty}} \sup_{x \in \bar{\mathbb{U}}} \left| \int_a^T [\mathbb{P}_x(X(t) \in \mathbb{U}_{1,n}) - \mathbb{P}_x(X(t) \in \mathbb{U}_{2,n})] dt - \int_a^T [p(t, x, \mathbb{U}_1) - p(t, x, \mathbb{U}_2)] dt \right| = 0. \quad (4.23)$$

Hence we have shown that the three terms which we decomposed $G_n(x)$ into at the beginning of the proof can be made arbitrarily close to the corresponding terms for $G(x)$ for large n . Thus the proof is complete. $\square$

Proof of Theorem 3.1. The proof follows immediately from Lemma 4.8 and Theorem 4.2. $\square$

5. PART II: HITTING MEASURE ESTIMATES ON $\mathbb{U}_n$

The first result is a Beurling type estimate which says that for any connected subset A of $\mathbb{U}_n$ with large enough diameter which is at a certain distance away from $\mathbb{U}_{1,n}$ the probability that random walk started from a neighboring site of A , hits $\mathbb{U}_{1,n}$ before hitting A decays as a power law in n . We first need the following definition.

Definition 3. Let $\mathbb{U}_n^*$ denote the graph $\mathbb{U}_n$ along with all the diagonals of the squares that are entirely in $\mathbb{U}_n$. We will call connected subsets of $\mathbb{U}_n^*$ as $*$ -connected subsets of $\mathbb{U}_n$.

Also recall the definition of random walk from (3.1). In the sequel for any subset $A \subset \mathbb{U}_n$, $\tau(A)$ will denote the hitting time for the random walk.

Lemma 5.1. Fix $c > 0$. Consider $A \subset \mathbb{U}_n$ be $*$ -connected. Also suppose that $\min(\text{diam}(A), d(\mathbb{U}_1, A)) \geq c$. Then for large n , for all such A ,

$$\sup_{x \sim A} \mathbb{P}_x(\tau(\mathbb{U}_{1,n}) \leq \tau(A)) \leq \frac{C}{n^\beta}$$

for some positive β, C depending only on c and $\mathbb{U}$. Here $x \sim A$ means that $x \notin A$ and there exists $y \in A$ such that x is a neighbor of y .

The next lemma says that uniformly from any point z at distance $1/2$ (any constant would work) from x_1 and all subsets $A \subset \mathbb{U}_n$ of large enough π_{RW} measure, the chance that the random walk does not hit A before reaching a ball of radius ϵ from x_1 goes to 0 as ϵ goes to 0.

Lemma 5.2. *Fix a constant $\alpha \in (0, 1)$. For all $\mathbb{U}, x_1$ as in Setup 1,*

$$\lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \sup_{\substack{A \subset \mathbb{U}_n \\ \pi_{RW}(A) \geq \alpha}} \left[\sup_{z \in \mathbb{U}_n \setminus B(x_1, \frac{1}{2})} \mathbb{P}_z \{ \tau(B(x_1, \epsilon)) \leq \tau(A) \} \right] = 0$$

where $\pi_{RW}(\cdot)$ is the stationary measure of the random walk on $\mathbb{U}_n$ and $B(x_1, \epsilon)$ is the euclidean ball of radius ϵ around x_1 .

The next lemma compares the hitting times of various sets i.e. how do the hitting times of various sets at different distances from the starting point compare.

Lemma 5.3. *Given small enough $\epsilon > 0$, for all $z \in \mathbb{U}_n \cap \mathbb{U}_{(\alpha - \sqrt{\epsilon})}$, and $y \in \mathbb{U}_n$ such that $d(y, z) \leq \epsilon^2$,*

$$\mathbb{P}_y(\tau(z) < \tau(\mathbb{U}_n \setminus \{\mathbb{U}_n \cap \mathbb{U}_{(\alpha)}\})) = \Theta\left(\frac{\log(1/d(y, z))}{\log n}\right),$$

where the constant in the Θ notation depend on $\epsilon, \alpha, \mathbb{U}$.

Remark 5.1. One can prove the above lemma from conductance estimates on the graph $\mathbb{U}_n$. However for the sake of unification, all the proofs in this article will use heat kernel estimates for the random walk on $\mathbb{U}_n$ which we state soon.

As mentioned before the above estimates are standard for the random walk on the entire lattice $\mathbb{Z}^2$. We make necessary adaptations to obtain the results for the random walk on the bounded geometry $\mathbb{U}_n$. We start by stating a basic mixing time result for the random walk on $\mathbb{U}_n$.

Lemma 5.4. *Given a smooth domain $\mathbb{U}$ as in Setup 1 and $\epsilon > 0$ there exists a constant $C = C(\epsilon, \mathbb{U})$ such that for large enough n ,*

$$t_\infty(\epsilon) \leq C$$

where $t_\infty(\cdot)$ is the $\mathbb{L}_\infty$ mixing time for the random walk on $\mathbb{U}_n$.

Proof. The fact is standard and the proof follows from [12, Theorem 13] and the isoperimetric inequality proved in [7, Theorem 5.5]. $\square$

We now state a standard property about the boundary of $\mathbb{U}$ as in Setup 1. Since the boundary $\partial\mathbb{U}$ is analytic there exists a $C > 0$ and an ϵ_0 such that for all $x \in \partial\mathbb{U}$ there exists an orthogonal system of coordinates centered at $x = (x_1, x_2)$ such that for all $\epsilon \leq \epsilon_0$

$$B(x, \epsilon) \cap \mathbb{U} = \{(x'_1, x'_2) \in B(x, \epsilon) : x'_1 \in (x_1 - \epsilon, x_1 + \epsilon), x'_2 \geq f(x'_1)\} \quad (5.1)$$

and

$$|f(x'_1) - x_2| \leq C|x'_1 - x_1|^2.$$

The above is a simple consequence of Taylor expansion up to second order of the curve locally near x . See Fig 2.

As a simple corollary of the above fact we see that $\mathbb{U}$ satisfies the following property which shows that the y_i 's in Setup 1 can indeed be chosen.




FIGURE 2. Locally the region near the boundary looks like a half plane

Corollary 5.1. *Let $\mathbb{U}$ be as in Setup 1. Then there exists $\delta_0 = \delta_0(\mathbb{U})$ such that for all $x \in \overline{\mathbb{U}}$ and $\delta < \delta_0$ there exists $y \in \mathbb{U}$ such that*

$$d(y, x) \leq \delta \quad (5.2)$$

$$B(y, \delta/2) \subset \mathbb{U}. \quad (5.3)$$

Recall that $d(y, x)$ is the euclidean distance between x and y . $B(y, \delta)$ denotes the euclidean ball of radius δ with center at y .

Proof. Choose $\delta_0 \leq \epsilon_0/4$ such that

$$C\delta_0^2 \leq \frac{\delta_0}{100}$$

where ϵ_0 and C appear in (5.1). For any $\delta < \delta_0$ the lemma is immediate if $d(x, \partial\mathbb{U}) > \delta/2$. since then we can choose $y = x$.

Otherwise let $z = (z_1, z_2) \in \partial\mathbb{U}$ be the closest point on the boundary to x . Now in the local coordinate system centered at z as in (5.1) choose $y = (z_1, z_2 + \delta/2)$. Then

$$d(x, y) \leq d(x, z) + d(z, y) \leq \delta.$$

Also clearly $B(y, \delta/2) \subset \mathbb{U}$ and hence we are done. See Fig 3. $\square$

For the remaining part we need the following gaussian upper and lower bounds on the heat kernel of random walk on $\mathbb{U}_n$. Recall the definition of $p_n(t, x, y)$ from (4.20).

Theorem 5.2. *Given $\mathbb{U}$ as in Setup 1,*

- i. [7, Theorem 2.9] *for any T there exists C_1 and C_2 such that for all $t \in [\frac{1}{n}, T]$,*

$$p_n(t, x, y) \leq \frac{C_1}{(t^{1/2} \vee \frac{1}{n})^2} \exp(-C_2 \frac{d(x, y)^2}{t}), \quad (5.4)$$

- ii. [8, Cor 2.2.5] *for any $T > 0$ there exists $C = C$ and N such that*

$$\mathbb{P}_x[\sup_{s \leq t} d(X_s, x) \geq \eta] \leq C \exp\left(t - \frac{\eta}{4(\frac{1}{n} \vee t^{1/2})}\right) \quad (5.5)$$

for all $n > N$, $t \leq T$, $x \in \mathbb{U}_n$, $\eta > 0$,

- iii. [7, Theorem 2.10] *there exists C_1, C_2 dependent only on T such that*

$$p_n(t, x, y) \geq \frac{C_1}{(\frac{1}{n} \vee t^{1/2})^2} \exp\left(-C_2 \frac{d(x, y)^2}{t}\right) \quad (5.6)$$

for all $t \leq T$ and $x, y \in \mathbb{U}_n$.

Note that all the constants above implicitly depend on $\mathbb{U}$.




FIGURE 3. Illustrating the proof of Corollary 5.1.

We now proceed towards proving Lemma 5.1. To this end we need a preliminary result. We start with a definition. Fix any $x \in \mathbb{U}_n$. Let us consider concentric discs $D_{x,j}$ around x of radius $\frac{2^j}{n}$ i.e.

$$D_{x,j} := B(x, \frac{2^j}{n}) \cap \mathbb{U}_n. \quad (5.7)$$

Recall ∂_{out} from (2.1). Let

$$C_{x,j} := \partial_{out} D_{x,j}. \quad (5.8)$$

We now show that starting from any point in $\mathbb{U}_n$ which is $\frac{1}{n}$ distance away from $C_{x,j-1}$ there is a constant chance of the random walk moving away from the boundary of $\mathbb{U}$, before hitting $C_{x,j}$. That is, it is unlikely that the random walk path on $\mathbb{U}_n$ hitting $C_{x,j}$ from $C_{x,j-1}$ stays uniformly close to the boundary of $\mathbb{U}$. The next result makes the above statement precise. Let $\epsilon < \delta_0$ where δ_0 appears in the statement of Corollary 5.1. Now by Corollary 5.1 for every $j \leq \log(\delta_0 n)$ and all $z \in C_{x,j-1}$, there exists a y such that

$$d(z, y) \leq \frac{\epsilon 2^{j-1}}{n}, \text{ and } B(y, \frac{\epsilon 2^{j-1}}{2n}) \subset \mathbb{U}.$$

Define,

$$B_{z,j-1} := B(y, \frac{\epsilon 2^{j-1}}{4n}). \quad (5.9)$$

For notational brevity we choose to suppress the ϵ dependence in the notation above.

Lemma 5.5. *There exists constants ϵ, c such that for all $x \in \mathbb{U}_n$, for any $z \in C_{x,j-1}$ with $j = 1, \dots, \log(\delta_0 n)$,*

$$\mathbb{P}_z(\tau(B_{z,j-1}) < \tau(C_{x,j})) \geq c,$$

where δ_0 appears in the statement of Corollary 5.1.

See Fig. 4.

Proof. To prove the lemma we use Theorem 5.2 ii. and iii. Since for all $z \in C_{x,j-1}$, $d(z, C_{x,j}) \geq \frac{2^{j-1}}{n}$ taking $T = 1$, $t = \epsilon^2 \frac{2^{2j}}{n^2}$ and $\eta = \frac{2^{j-1}}{n}$, by (5.5) we have

$$\mathbb{P}_z[\tau(C_{x,j}) \leq t] \leq Ce \exp\left(-\frac{1}{4\epsilon}\right).$$



FIGURE 4. The left hand side shows the unlikely event that a path from $C_{x,j-1}$ hits $C_{x,j}$ and the path stays close to the boundary of $\mathbb{U}$. The right hand side shows a typical path.

Now we look at the chance that $\tau(B_{z,j-1}) \leq t$. Clearly

$$\mathbb{P}_z(\tau(B_{z,j-1}) \leq t) \geq \mathbb{P}_z(X(t) \in B_{z,j-1}).$$

Let us recall from (4.20) that by definition

$$\mathbb{P}_z(X(t) \in B_{z,j-1}) = \sum_{y \in B_{z,j-1}} p_n(t, z, y) m_n(y).$$

Now since all the points in $B_{z,j-1} \cap \mathbb{U}_n$ have 4 neighbors $m_n(y) = \frac{1}{n^2}$ for all $y \in B_{z,j-1}$. Let us for the moment denote it by m . By (5.6) we get

$$\mathbb{P}_z(X(t) \in B_{z,j-1}) \geq C_1 m \frac{|B_{z,j-1} \cap \mathbb{U}_n|}{t} \exp(-C_2) = \frac{\epsilon^2 2^{2(j-1)}}{\pi n^2 \epsilon^2 2^{2j} / n^2} = \Theta(1).$$

Thus

$$\begin{aligned} \mathbb{P}_z(\tau(B_{z,j-1}) < \tau(C_{x,j})) &\geq \mathbb{P}_z(\tau(B_{z,j}) < t) - \mathbb{P}_z(\tau(C_{x,j}) < t) \\ &\geq \Theta(1) - O(\exp(-\frac{1}{4\epsilon})). \end{aligned}$$

Thus we are done by choosing ϵ small enough. $\square$

We are now ready to prove Lemma 5.1. As remarked earlier the basic structure of the arguments are standard and are used to prove similar statements on the whole lattice. We make the necessary additional arguments to prove the statement in the bounded geometry of $\mathbb{U}_n$.

Proof of Lemma 5.1. We will fix $\delta_1 < \delta_0/4 \wedge c$ where δ_0 appears in Corollary 5.1 and c appears in the statement of the lemma. Let $x \sim A$. Recall the definitions of $D_{x,j}$ and $C_{x,j}$ from (5.7) and (5.8) with center x . Let us denote by

$$I_j := D_{x,j} \setminus D_{x,j-2}.$$

for $j = 1 \dots \log n \delta_1$. Now given any x if $d(x, \partial\mathbb{U}) \geq \delta_1$ then for all j , I_j is an annulus. Otherwise some of the I_j 's are a topological quadrilateral and two of the four sides are a part of $\partial\mathbb{U}$. See Fig




FIGURE 5. Random walk completing a full circle oriented in one direction will hit the set A .

5 ii. If δ_1 is chosen to be small enough clearly these are the only two possibilities. For $z \in C_{x,j}$ recall $B_{z,j}$ from (5.9). By Lemma 5.5 there is a positive c_1 such that for all $j = 1, \dots, \log(\delta_1 n)$,

$$\inf_{z \in C_{x,j}} \mathbb{P}_z(\tau(B_{z,j}) \leq \tau(C_{x,j+1})) \geq c_1. \quad (5.10)$$

Now any point in $B_{z,j}$ is in the interior of I_j and at distance at least $\frac{\epsilon 2^{j-1}}{4n}$ from $C_{x,j+1} \cup C_{x,j-1} \cup \partial \mathbb{U}$. Let z' be any such point. Since by hypothesis A is a $*$ -connected set and $\text{diam}(A) > c$ the following observations are straightforward corollaries of the Jordan curve theorem: for $j = 1 \dots \log(\delta_1 n)$,

- If I_j is an annulus: any curve starting from z' which stays $\frac{\epsilon}{100} \frac{2^{j-1}}{n}$ away from the boundary, makes a full circle and completes a closed loop hits A . Fig 5 i.
- If I_j is a topological quadrilateral: Out of the four sides, two sides of I_j are a part of $\partial \mathbb{U}$. There exists one of these two sides such that any curve γ which starts from z' and stays $\frac{\epsilon}{100} \frac{2^{j-1}}{n}$ away from the three sides before hitting that side hits A . Fig 5 ii.

By the Donsker invariance principle, simple random walk does both the above things with constant probability only dependent on ϵ and independent of j . Thus the chance that random walk started from x hits A between $\tau(C_{x,j})$ and $\tau(C_{x,j+1})$ is at least d for some constant $d = d(\epsilon)$. This follows since from the location z at time $\tau(C_{x,j})$ the random walk with constant chance hits $B_{z,j}$ and from there hits A with constant probability before $\tau(C_{x,j+1})$.

We consider the time interval $[\tau(C_{x,j}), \tau(C_{x,j+1}))$ as the j^{th} round. By the previous discussion in each round the chance to hit A is at least d for $j = 1 \dots \log(n\delta_1)$. Also since by hypothesis

$d(A, \mathbb{U}_1) \geq c$ and $x \sim A$,

$$\tau(C_{x,j}) < \tau(\mathbb{U}_1)$$

for all such j . Hence

$$\sup_{x \sim A} \mathbb{P}_x(\tau(\mathbb{U}_1) \leq \tau(A)) \leq (1-d)^{\log(\delta_1 n)}.$$

□

We now state and prove another similar lemma. Recall the definitions of $y_1, \mathbb{U}_{1,n}$ from Setup 1.

Lemma 5.6. *Let $0 < \epsilon_1 < \epsilon_2$. Assume $A \subset \mathbb{U}_n$ is a connected set such that*

$$d(\mathbb{U}_{1,n}, A) \leq \epsilon_1.$$

Also assume $A \cap (\mathbb{U}_n \setminus B(y_1, \epsilon_2)) \neq \emptyset$. Then

$$\sup_{x \in \mathbb{U}_{1,n}} \mathbb{P}_x(\tau(\mathbb{U}_n \setminus B(y_1, \epsilon_2)) \leq \tau(A)) \leq C^{\log(\frac{\epsilon_2}{\epsilon_1})},$$

for some $C = C(\mathbb{U}) < 1$ independent of n .




FIGURE 6. Figure illustrating the proof of Lemma 5.6. The red random walk path making a loop hits A .

The statement of the lemma roughly says if a connected set A of large enough diameter is close enough to $\mathbb{U}_{1,n}$ then random walk starting from $\mathbb{U}_{1,n}$ is more likely to hit the set A before exiting a large enough ball.

Proof. The proof of this lemma is similar to the proof of Lemma 5.1. We look at shells of exponentially growing radii centered at y_1 i.e. $B(y_1, 2^j \epsilon_1) \cap \mathbb{U}$ for $j = 1 \dots \log(\frac{\epsilon_2}{\epsilon_1})$. Recall ∂_{out} from (2.1). Let

$$C_j = \partial^{out}(B(y_1, 2^j \epsilon_1) \cap \mathbb{U}_n).$$

Let $\tau(C_j)$ be the first time that the random walk hits C_j . We first claim that there is a constant c such that for all j ,

$$\inf_{z \in C_j} \mathbb{P}_z(\tau(A) \leq \tau(C_{j+1})) \geq c.$$

The proof of the above claim is the same as the proof of Lemma 5.1. We omit the arguments to avoid repetition.

Now let the random walk start anywhere from $\mathbb{U}_{1,n}$ and let $z \in C_j$ be the point it hits at $\tau(C_j)$. Thus again using the round argument as in the proof of the previous lemma

$$\sup_{x \in \mathbb{U}_{1,n}} \mathbb{P}_x(\tau(\mathbb{U} \setminus B(x_1, \epsilon_2)) \leq \tau_A) \leq (1 - c)^{\log \frac{\epsilon_2}{\epsilon_1}}.$$

Note that the number of rounds here is $\log \frac{\epsilon_2}{\epsilon_1}$. □

Proof of Lemma 5.2. We first prove the following lemma:

Lemma 5.7. *Fix $\alpha \in (0, 1)$. Then there exists constants C, D such that for all large enough t ,*

$$\sup_{\substack{A \subset \mathbb{U}_n \\ \pi_{RW}(A) \geq \alpha}} \sup_{x \in \mathbb{U}_n} \mathbb{P}_x(\tau(A) \geq t) \leq Ce^{-Dt}$$

where $\pi_{RW}(\cdot)$ is the stationary measure of the random walk on $\mathbb{U}_n$.

This is a standard mixing result which says that the hitting time of any set A of large enough measure, for the random walk has exponential tail.

Proof. Fix any $A \subset \mathbb{U}_n$. As stated in Lemma 5.4 $\mathbf{t} := t_\infty(\frac{1}{4}) = O(1)$. Now since $\pi_{RW}(A) \geq \alpha$, $\inf_{x \in \mathbb{U}_n} \mathbb{P}_x(X_{\mathbf{t}} \in A) \geq \alpha/4$, and hence

$$\inf_{x \in \mathbb{U}_n} \mathbb{P}_x(\tau(A) \leq \mathbf{t}) \geq \alpha/4.$$

Therefore for any $t > 0$

$$\mathbb{P}_x(\tau(A) \geq t) \leq (1 - \alpha/4)^{\lfloor t/\mathbf{t} \rfloor}.$$

This is because from any $y \in \mathbb{U}_n$ there is a chance of at least $\alpha/4$ to hit B_1 in the next time interval of length $\mathbf{t}$. Hence the lemma is proved. □

We resume the proof of Lemma 5.2. To prove this we compute the time spent in $B(x_1, \epsilon)$ starting from a point z_1 and z_2 such that $d(x, z_1) \leq 2\epsilon$ and $d(x, z_2) > \frac{1}{2}$. This is helpful because of the following formula:

$$\mathbb{P}_{z_2}(\tau(B(x_1, \epsilon)) < T) \leq \frac{\int_0^{2T} \mathbb{P}_{z_2}(t, B(x_1, \epsilon))}{\inf_{z_1} \int_0^T \mathbb{P}_{z_1}(t, B(x_1, \epsilon))}. \quad (5.11)$$

The above follows by markov property. The goal now is to prove upper and lower bounds on the numerator and the denominator respectively. We use the upper and lower bound on the gaussian

heat kernel stated in Theorem 5.2. Assume $T \geq 1$. We see that

$$\begin{aligned} \int_0^T \mathbb{P}_{z_1}(t, B(x_1, \epsilon)) &\geq \int_\epsilon^T \left[\sum_{y \in B(x_1, \epsilon)} \frac{1}{n^2} p_n(t, z_1, y) \right] dt \\ &\geq \int_\epsilon^1 \left[\sum_{y \in B(x_1, \epsilon)} \frac{1}{n^2} p_n(t, z_1, y) \right] dt \\ &\geq C\epsilon^2 \int_\epsilon^1 \frac{1}{t} dt \\ &\geq C\epsilon^2 \log\left(\frac{1}{\epsilon}\right). \end{aligned}$$

The second last inequality follows from (5.6) (the constant C is changing from line to line). We now look at the numerator. Recall $\mathbf{t}$ from the proof of the last lemma.

$$\begin{aligned} \int_0^{2T} \mathbb{P}_{z_2}(t, B(x_1, \epsilon)) &\leq \epsilon P_{z_2}(\tau(B(x_1, \epsilon)) \leq \epsilon) + \int_\epsilon^{2T} \left[\sum_{y \in B(x_1, \epsilon)} \frac{1}{n^2} p_n(t, z_2, y) \right] dt \\ &= \epsilon P_{z_2}(\tau(B(x_1, \epsilon)) \leq \epsilon) + \int_\epsilon^{\mathbf{t}} \left[\sum_{y \in B(x_1, \epsilon)} \frac{1}{n^2} p_n(t, z_2, y) \right] dt + \int_{\mathbf{t}}^{2T} \left[\sum_{y \in B(x_1, \epsilon)} \frac{1}{n^2} p_n(t, z_2, y) \right] dt \\ &\leq \epsilon O(e^{-\frac{C}{\sqrt{\epsilon}}}) + C\epsilon^2 \int_\epsilon^{\mathbf{t}} \frac{1}{t} e^{-\frac{C}{t}} dt + C\epsilon^2 T \\ &\leq CT\epsilon^2. \end{aligned}$$

The bounds on the first and second terms follow from (5.5) and (5.4) respectively. Note for the last integral we use the fact that for any $t \geq \mathbf{t}$, $p_n(t, z_2, y) \leq \frac{C}{n^2}$. Thus using (5.11)

$$\mathbb{P}_{z_2}(\tau(B(x_1, \epsilon)) < T) \leq O\left(\frac{T}{\log(\frac{1}{\epsilon})}\right).$$

Now by Lemma 5.7

$$\mathbb{P}_{z_1}\{\tau(A) < T\} \geq 1 - O(e^{-DT}).$$

Using the above bounds, taking $z = z_2$ we get that for any $T \geq 1$

$$\begin{aligned} \mathbb{P}_z\{\tau(B(x_1, \epsilon)) \leq \tau(A)\} &\leq \mathbb{P}_z\{\tau(B(x_1, \epsilon)) < T\} + \mathbb{P}_z\{\tau(A) \geq T\} \\ &\leq O\left(\frac{T}{\log(\frac{1}{\epsilon})}\right) + O(e^{-DT}). \end{aligned}$$

Thus we are done by choosing $T = \sqrt{\log(\frac{1}{\epsilon})}$. □

Proof of Lemma 5.3. The proof uses similar arguments as above. We begin by providing sharp upper and lower bounds for the two following quantities:

$$\begin{aligned} \mathbb{E}_y\left[\int_0^\tau 1(X(t) = z)\right] \\ \mathbb{E}_z\left[\int_0^\tau 1(X(t) = z)\right] \end{aligned}$$

where $\tau = \tau(\mathbb{U}_n \cap [\mathbb{U} \setminus \mathbb{U}_{(\alpha)}])$. To show this we first notice that by either Lemma 5.4 or Theorem 5.2 iii., there exists a constant $c = c(\mathbb{U}, \alpha)$ for any $w \in \mathbb{U}_n$,

$$\mathbb{P}_w(\tau \leq 1) \geq c.$$

Using Theorem 5.2 we get for any $w \in \mathbb{U}_n$,

$$\int_0^2 \mathbb{P}_w(X(t) = z) \leq \frac{1}{n^2} \int_0^2 \frac{e^{-\frac{d(w,z)^2}{t}}}{\frac{1}{n^2} \vee t} dt = O\left(\frac{\log(1/d(w,z))}{n^2}\right). \quad (5.12)$$

Putting the above together,

$$\begin{aligned} \mathbb{E}_w\left[\int_0^\tau 1(X(t) = z)\right] &\leq \int_0^2 \mathbb{P}_w(X(t) = z) + \sum_{\ell=0}^\infty \mathbb{P}_w(\tau \geq \ell) \sup_{w \in \mathbb{U}_n} \int_{\ell+1}^{\ell+2} \mathbb{P}_w(X(t) = z) \\ &\leq O\left(\frac{\log(1/d(w,z))}{n^2}\right) + \sum_{\ell=0}^\infty (1-c)^\ell \sup_{w \in \mathbb{U}_n} \int_1^2 \mathbb{P}_w(X(t) = z) \\ &= O\left(\frac{\log(1/d(w,z))}{n^2}\right) + O\left(\frac{1}{n^2}\right). \end{aligned}$$

The second term follows by the fact that starting from any w in time 1 the random walk is reasonably mixed (by Theorem 5.2) and then from the location of the random walk at time 1 we use (5.12). For the lower bound we use strong Markov Property,

$$\mathbb{E}_y\left[\int_0^\tau 1(X(t) = z)\right] \geq \int_0^{\epsilon^2} \mathbb{P}_y(X(t) = z) - \mathbb{P}_y(\tau \leq \epsilon^2) \left[\sup_{w: d(w,z) > \sqrt{\epsilon}} \int_0^{\epsilon^2} \mathbb{P}_w(X(t) = z) \right]$$

Now for any $y \in \mathbb{U}_n$ such that $d(y, z) \leq \epsilon^2$,

$$\begin{aligned} \int_0^{\epsilon^2} \mathbb{P}_y(X(t) = z) &\geq \frac{1}{n^2} \int_0^{\epsilon^2} \frac{1}{\frac{1}{n^2} \vee t} e^{-\frac{d(y,z)^2}{t}} dt \\ &\geq \frac{1}{n^2} \int_{d(y,z)^2}^{\epsilon^2} \frac{1}{t} e^{-1} dt \\ &\geq \Omega\left(\frac{\log(1/(d(y,z)))}{n^2}\right) \end{aligned}$$

where the last inequality follows since $d(y, z) \leq \epsilon^2$ by hypothesis.

By Theorem 5.2, $\mathbb{P}_y(\tau \leq \epsilon^2) \leq O(e^{-\frac{1}{\sqrt{\epsilon}}})$. Also for any w such that $d(w, z) \geq \sqrt{\epsilon}$ we have

$$\int_0^{\epsilon^2} \mathbb{P}_w(X(t) = z) \leq \frac{1}{n^2} \int_0^{1/n^2} \frac{1}{\frac{1}{n^2}} + \frac{1}{n^2} \int_0^{\epsilon^2} \frac{e^{-\frac{\epsilon}{t}}}{t} dt$$

Thus from the above we see that for all y such that $d(y, z) \leq \epsilon^2$ we have,

$$P_y(\tau(z) < \tau(\mathbb{U} \setminus \mathbb{U}_\alpha)) = \frac{\int_0^\tau P_y(X(t) = z)}{\int_0^\tau P_z(X(t) = z)} = \frac{1}{\log(n)} \Theta(\log(1/d(x, z))),$$

and we are done. $\square$

REFERENCES

- [1] Lars V Ahlfors. *Complex analysis: an introduction to the theory of analytic functions of one complex variable*. 1979.
- [2] Richard F Bass. Brownian motion, heat kernels, and harmonic functions. In *Proceedings of the International Congress of Mathematicians*, pages 980–985. Springer, 1995.
- [3] Richard F Bass and Pei Hsu. Some potential theory for reflecting Brownian motion in holder and lipschitz domains. *Annals of Probability*, pages 486–508, 1991.
- [4] Heinrich Begehr. Boundary value problems in complex analysis ii. *Boletín de la Asociación Matemática Venezolana*, page 217, 2005.
- [5] Krzysztof Burdzy and Zhen-Qing Chen. Discrete approximations to reflected Brownian motion. *Annals of Probability*, 36(2):698–727, 2008.
- [6] Zhen-Qing Chen. On reflecting diffusion processes and skorokhod decompositions. *Probability theory and related fields*, 94(3):281–315, 1993.
- [7] Zhen-Qing Chen and Wai-Tong Fan. Hydrodynamic limits and propagation of chaos for interacting random walks in domains. *arXiv preprint arXiv:1311.2325*, 2013.
- [8] Wai-Tong Fan. *Interacting particle systems with partial annihilation through membranes*. PhD thesis, University of Washington, 2014.
- [9] S Ganguly and Y Peres. Competitive erosion is conformally invariant. *Manuscript*, 2015.
- [10] Harry Kesten. Relations between solutions to a discrete and continuous dirichlet problem. In *Random walks, Brownian motion, and interacting particle systems*, pages 309–321. Springer, 1991.
- [11] David Asher Levin, Yuval Peres, and Elizabeth Lee Wilmer. *Markov chains and mixing times*. American Mathematical Soc., 2009.
- [12] Ben Morris and Yuval Peres. Evolving sets, mixing and heat kernel bounds. *Probability Theory and Related Fields*, 133(2):245–266, 2005.
- [13] Christian Pommerenke. *Boundary behaviour of conformal maps*. Springer, 1992.
- [14] Ken-iti Sato and Tadashi Ueno. Multi-dimensional diffusion and the markov process on the boundary. *Journal of Mathematics of Kyoto University*, 4(3):529–605, 1965.
- [15] Stanislav Smirnov. Discrete complex analysis and probability. *arXiv preprint arXiv:1009.6077*, 2010.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, WA, USA.

E-mail address: `sganguly@math.washington.edu`

MICROSOFT RESEARCH, WA, USA.

E-mail address: `peres@microsoft.com`